ALLAMA IQBAL OPEN UNIVERSITY

USMAN NOTES
www.usmannotes.com

Name:

Registration No.:

Course Code: 2511

Program: MA/MSc/ M.Ed

Semester: Autumn 2026

ALLAMA IQBAL OPEN UNIVERSITY, ISLAMABAD

(Department of Agricultural Science)

Course: Soil Conservation & Watershed Management (2511)

Level: MA/MSc/ M.Ed

Semester: Autumn 2026

Assignment No. 2 (Units 5-9)

Total Marks: 100 • Pass Marks: 50

SOLVED ASSIGNMENT GUIDE — For Study & Reference

Note: This guide is prepared for study and reference purposes only. Students are advised to read the relevant units of the course book, understand the concepts, and then write the assignment in their own words. Do not copy this guide word for word.

USMAN NOTES www.usmannotes.com

Question 1: Causes and Impacts of Land Degradation (20 Marks)

Q. 1 Explain the causes and impacts of land degradation. (20)

Introduction

Land degradation is the decline in the productive capacity and ecological functioning of land — the process by which soil, vegetation and water resources deteriorate so that the land yields less, supports less life and becomes harder and more expensive to restore the longer the process continues. It is not a single problem but a family of processes, including soil erosion by water and wind, loss of soil organic matter and nutrients, salinisation and waterlogging, compaction, the decline of vegetation cover and the loss of biodiversity, often acting together

on the same field. In Pakistan the problem touches every major land system: the eroded rainfed tracts of the Potohar and the northern hills, the wind-swept arid zones, the salt-affected irrigated plains and the overgrazed rangelands that cover much of the country's area. This answer first explains the causes of land degradation — distinguishing the natural background, the direct human activities that trigger it, and the underlying social and economic drivers — and then describes its impacts on production, on water and the environment, and on the incomes and welfare of the people who depend on the land.

Direct Causes: Unsustainable Land Use and Management

The immediate causes of degradation are the ways land is used and managed beyond its capacity. Cultivation of steep and marginal land, ploughing up and down the slope, repeated tillage that pulverises structure, and the removal or burning of crop residues expose soil to water and wind erosion, while continuous cropping without adequate return of manure, fertiliser or fallow mines the soil's nutrients and organic matter until structure collapses and yields fall even where erosion is not visible. Overgrazing is the dominant direct cause on rangelands and common lands: too many animals for too long remove the protective cover, select out the palatable species, compact the surface with trampling and leave the soil open to both runoff and wind, so the pasture degrades in a spiral in which poorer cover supports still poorer regrowth. Deforestation and unsustainable cutting for fuelwood, timber and land clearing strip slopes and catchments of protection, starting gullies and raising sediment loads. In irrigated areas the characteristic direct causes are different: over-irrigation and poor drainage raise the water table and bring salts into the root zone, producing waterlogging and salinisation; irrigation with poor-quality groundwater adds sodium that disperses soil structure;

and the neglect of field levelling and drainage maintenance spreads the damage. To these must be added urbanisation, mining, road building and waste disposal, which seal, excavate or contaminate soil outright. In every case the common feature is use that takes more from the land system — soil depth, nutrients, cover, drainage — than the management returns to it.

Underlying Causes: Population, Poverty, Tenure and Policy

Direct causes persist because underlying conditions drive people toward them. Population growth subdivides holdings, pushes cultivation onto steeper and drier margins, multiplies livestock on a fixed grazing resource and creates the fuelwood demand that strips hillsides; poverty then compels short-term survival decisions, because a family that must eat this season cannot easily fallow, terrace or replant for a benefit ten years away. Insecure land tenure discourages investment in protection — tenants and sharecroppers who may lose the plot have little reason to build bunds or plant trees on it — while the open-access character of much grazing land removes any individual's incentive to rest it, the classic tragedy of the commons. Market and policy signals can deepen the problem: prices and subsidies that reward water-intensive cropping in dry areas, the underpricing of irrigation water and energy for pumping, weak extension advice on conservation, and the historic treatment of 'wasteland' development as simply bringing more area under the plough all encourage degradation unintentionally. Climate change now acts as a force multiplier on all of these, intensifying the storms that erode, lengthening the droughts that bare the soil and shifting the conditions to which existing farming systems were adapted. A serious answer about causes must hold both levels together: blaming only the farmer's practice misses the pressures that make the practice rational, and blaming

only the pressures misses the practical changes in management through which recovery actually begins.




Figure 1: Common forms and drivers of land degradation (illustrative severity index — a teaching diagram, not measured data).

Process Forms of Degradation

It helps the discussion of impacts to name the main forms degradation takes. Water erosion removes topsoil selectively, carrying away the fine, fertile fraction first and, in its gully form, physically consuming fields, roads and settlements. Wind erosion does the same selective removal in dry regions and buries neighbouring land and canals under the coarser material it drops. Chemical degradation includes nutrient depletion, acidification in some environments, pollution, and the salinisation and sodification that are the leading chemical problems in irrigated Pakistan. Physical degradation covers compaction by machinery and hooves, crusting and sealing of the surface, loss of structure and the waterlogging that drowns soil air. Biological degradation — the decline of soil organic matter, soil organisms and above-ground biodiversity

— underlies and accelerates all the others, because a biologically impoverished soil infiltrates less water, holds fewer nutrients and resists erosion less. Most degraded land suffers several forms at once, which is why single-measure reclamation so often disappoints and why the integrated practices discussed later in this assignment are necessary.

Impacts on Agricultural Production and Food Security

The first impact is on production itself, on-site where the degradation occurs. Loss of topsoil means loss of rooting depth, stored water and nutrient capital, so yields fall and become more variable, responding less to fertiliser because the soil can no longer hold and supply it efficiently; farmers compensate with higher input doses at higher cost per unit of output, or abandon the worst land altogether. Salinised and waterlogged fields drop out of production entirely in their severe patches, and gullied land is lost absolutely. At national scale these on-site losses add up to a widening gap between food demand and the productive capacity of a fixed or shrinking good-land base, increasing import dependence and exposure to world prices. Degradation also narrows future options: topsoil forms over centuries, so losses within a human generation are, for practical purposes, irreversible, and each hectare abandoned in the hills pushes cultivation pressure onto the remaining land, spreading the process.

Impacts on Water, Infrastructure and the Wider Environment

The second set of impacts falls off-site, on people who may live far from the degraded land. Sediment eroded from degraded catchments fills reservoirs and shortens their designed life, silts canals and raises their maintenance cost, chokes drainage systems, buries fertile valley fields under infertile sand and gravel, and raises river beds so that floods spread wider. The loss

of infiltration on degraded uplands converts steady streamflow into destructive flash floods followed by dry channels, damaging roads, bridges and settlements and reducing the groundwater recharge on which wells and springs depend. Dust from wind-eroded land harms health, transport and machinery. Environmentally, degradation releases stored soil carbon to the atmosphere, destroys habitat and native species, reduces the pollinators and natural pest enemies on which farming partly depends, and degrades the water quality of streams through sediment, nutrient and pesticide loads. These off-site costs are frequently larger than the on-site yield loss, and because they fall on the public rather than on the land user who caused them, they are the standard economic justification for public investment in conservation and watershed programmes.

Impacts on Livelihoods and Society

The third set of impacts is social. Degradation reduces and destabilises the incomes of farm and pastoral households, falling hardest on the poor, who hold the most marginal land, depend most on common grazing and fuel sources, and have the least capacity to buy their way out with inputs or to migrate for work; women bear a particular share of the burden, walking further for fuelwood and water as local resources decline. Falling land productivity fuels distress migration, disputes over shrinking good land and water, and the abandonment of villages in severely eroded tracts. It also destroys cultural and option values — local crop varieties, medicinal plants and landscapes — that markets do not price but communities feel. Conversely, because degradation is largely human-caused, it is also substantially reversible at reasonable cost where cover is restored, grazing is managed and soil organic matter is rebuilt,

as the remaining questions of this assignment describe; the impacts are therefore best read not as fate but as the price of delayed action, rising every year the causes are left to operate.

Table 1: Causes and impacts of land degradation — summary

Category	Main Items	Illustration
Direct causes	Erosive cultivation, residue removal, nutrient mining, overgrazing, deforestation, over-irrigation without drainage	Bare steep fields gully; salt patches spread in poorly drained canal commands
Underlying causes	Population pressure, poverty, insecure tenure, open-access commons, perverse prices and weak extension, climate change	Tenants do not terrace land they may lose; commons are grazed without rest
Production impacts	Yield decline and variability, rising input cost, land abandonment	Thin topsoil responds poorly to fertiliser; gullied plots leave cultivation
Off-site impacts	Reservoir and canal siltation, floods, poor recharge, dust, carbon release	Sediment from degraded catchments shortens storage life downstream

USMAN NOTES www.usmannotes.com

Social impacts	Income loss, fuel and water drudgery, migration, disputes — falling hardest on the poor	Longer daily walks for fuelwood as nearby cover disappears

Conclusion

In conclusion, land degradation is caused directly by land use and management that exceed the land's capacity — erosive cultivation, cover removal, nutrient mining, overgrazing, deforestation and irrigation without drainage — and indirectly by the population pressure, poverty, insecure tenure, open-access grazing, distorted incentives and intensifying climate that drive those practices. Its impacts run from falling and unstable yields and abandoned land on the farm, through silted reservoirs, floods, poor recharge and lost biodiversity in the wider environment, to reduced incomes, heavier drudgery and migration in society, with the poor bearing the heaviest share. Understanding both the direct and the underlying causes is essential, because effective remedies must combine better practices on the field with the grazing agreements, tenure security, incentives and community organisation that make those practices possible — the theme developed in the questions that follow.

Question 2: Agronomic Measures for Soil Conservation (20 Marks)

Q. 2 Discuss agronomic measures for soil conservation. (20)

Introduction

Agronomic measures for soil conservation are the methods that control erosion and build soil health through the choice, arrangement and management of crops and vegetation and through the handling of the soil surface — as distinct from structural measures, which reshape the land with earthworks. They work with the biology of the farming system itself: keeping living or dead cover on the soil for as much of the year as possible, improving the organic matter and structure that make soil resist detachment and absorb rain, arranging crops across the slope so that runoff is checked, and rotating and mixing crops so that fertility is maintained without exhausting the land. Their outstanding advantages are low cost, use of the farmer's existing skills and equipment, and the fact that they usually raise yields and income at the same time as they conserve, so that conservation is not a separate expense but a by-product of good farming. This answer discusses the main agronomic measures — vegetative cover and mulching, contour cultivation, strip cropping, crop rotation, mixed and intercropping, cover crops and green manuring, conservation tillage, and agroforestry and grass management in their agronomic aspects — and concludes with their limitations and place in an integrated programme.

Vegetative Cover, Mulching and Residue Management

The single most effective agronomic measure is simply to keep the soil covered. A living canopy intercepts raindrops before they can shatter aggregates, stems and litter slow surface flow so that sediment settles and water infiltrates, and roots hold the soil and maintain its pores. Where a living crop cannot occupy the ground — between rows, after harvest, before the next

sowing — dead cover performs much of the same work: retaining crop residues and stubble on the surface instead of burning or removing them, and spreading mulch of straw, stalks or other organic material on vulnerable plots, protects against both raindrop impact and wind lift, conserves soil moisture by shading the surface, moderates soil temperature and, as it decays, feeds the organisms that build stable structure. Residue burning, by contrast, destroys this protection at exactly the season when storms may strike, and deprives the soil of its main renewable source of organic matter. The practical target used in conservation teaching is continuous or near-continuous cover, planned through the cropping calendar so that bare, loose soil coincides as little as possible with the season of intense rain, choosing fast-establishing varieties and close spacing where early cover is needed. On rangeland and pasture the corresponding measure is managed grazing — stocking to the carrying capacity, rotational resting of paddocks, and reseeding of denuded patches — so that a protective grass cover is an actively maintained asset rather than an accident.

Contour Cultivation and Strip Cropping

Two agronomic measures rearrange field operations and crop layout across the slope. Contour cultivation, discussed fully in Assignment 1, performs ploughing, sowing and interculture on the level across the slope so that every furrow stores water instead of draining it; it costs nothing in materials and is the baseline practice for all sloping cropland. Strip cropping develops the same principle into a cropping pattern: the slope is divided into contour strips alternating between erosion-permitting crops, such as clean-tilled row crops, and erosion-resisting close-growing crops, such as grasses, legumes and dense cereals. Runoff leaving a cultivated strip enters the dense strip below at low energy, drops its sediment there — progressively building a natural

terrace effect as soil accumulates at strip boundaries — and infiltrates instead of gathering into rills. Variations include field strips of roughly uniform width on gentle, regular slopes, contour strips following the true contour on more critical land, buffer strips of permanent grass or shrubs protecting waterways and steep patches, and wind strips oriented across the prevailing wind in dry areas. Strip cropping conserves soil and moisture, diversifies production and risk on the same field, and can be adopted progressively without surrendering the whole holding to one system in a single year.

Crop Rotation, Mixed Cropping and Intercropping

What is grown in sequence and in mixture matters as much as how the surface is worked. A sound rotation alternates crops with different rooting depths, nutrient demands and canopy cover, and crucially includes a restorative phase — a legume, a fodder or a grass ley — that returns nitrogen and organic matter, rebuilds aggregate stability and breaks the pest cycles that force heavy tillage or chemical use. Rotating a soil-building crop before an erosion-prone but valuable row crop means the vulnerable phase is grown on soil in its most resistant condition. Mixed cropping and intercropping carry protection into the same season: a main crop is grown together with a companion — classically a cereal with a legume, or rows of a tall crop with a low, dense cover between them — so that ground cover is fuller and earlier, raindrops meet a layered canopy, roots occupy different soil layers, and the failure risk of the whole field is reduced because the components respond differently to drought or pest attack. Relay planting, in which the next crop is sown into the standing previous crop, shortens the bare interval between crops almost to nothing. These practices are traditional in much of Pakistan's rainfed farming, and conservation teaching builds on them by adjusting proportions, spacing and

species choice toward maximum cover and fertility return rather than replacing the farmer's system wholesale.

Cover Crops, Green Manuring and Fertility Management

Cover crops are grown specifically to protect and improve soil during a period when the land would otherwise be bare or between main crops: quick growing legumes and grasses sown after harvest, in orchards, or ahead of a main crop keep living roots in the ground, capture nitrate that would leach away, suppress weeds without tillage and add biomass. Green manuring ploughs such a crop into the soil in place, delivering organic matter exactly where it is needed with no transport cost. Around these practices sits the broader agronomy of soil fertility that conservation depends on: balanced use of manure, compost and fertiliser to maintain the nutrient supply that good cover growth requires; returning all available organic wastes to the land; liming or gypsum and salt management where soil chemistry limits plant growth, as in sodic soils; and building soil organic matter as the master variable, because organic matter simultaneously improves aggregate stability, infiltration, water holding and nutrient retention — the four soil properties that decide both erodibility and productivity. A conservation plan that ignores fertility will fail agronomically, since starved crops make thin cover; a fertility plan that ignores cover will fail physically, since well-fed soil still washes away if it is bare when the storm arrives.

Conservation Tillage and Agroforestry Elements

Conservation tillage reduces the disturbance that leaves soil naked and pulverised. Minimum tillage works the soil only where seed must be placed, and zero or no-till sowing places seed

directly into the residues of the previous crop, so the surface armour of stubble is never destroyed and the pore structure built by roots and worms is preserved; where herbicide dependence is a concern, cover-crop suppression and mechanical methods can substitute in part, and even a simple shift — fewer, shallower workings, done on the contour, avoiding tillage when the soil is wet enough to compact or dry enough to powder — captures much of the benefit with conventional equipment. Agroforestry contributes agronomic measures within farmland as well as the watershed role treated earlier: contour hedgerows of leguminous shrubs pruned to provide mulch and green manure while their roots hold the slope, trees on bunds and boundaries that recycle deep nutrients into litter, and fodder banks that feed livestock from a protected plot instead of from the open hillside. Grass management for conservation — contour grass strips, protected waterways, pasture improvement — completes the set by ensuring that the non-cropped parts of the farm also carry their protective cover.

USMAN NOTES www.usmannotes.com

Table 2: Agronomic measures for soil conservation — summary

Measure	How It Conserves	Additional Benefit
Cover and mulch / residues	Shield from raindrop and wind; slows runoff; feeds soil life	Moisture conservation; organic-matter build-up
Contour cultivation	Level furrows store water across the slope	Better moisture distribution; cheaper field work
Strip cropping	Dense strips trap sediment	Diversified output; gradual

	from cultivated strips	natural terracing
Rotation, mixed and intercropping	Restorative phases rebuild structure and nitrogen; fuller canopy	Yield stability; pest and risk reduction
Cover crops and green manuring	Living cover in bare periods; biomass returned in place	Nitrogen, weed suppression, nitrate capture
Fertility and organic-matter management	Stable aggregates; high infiltration and water holding	Higher, more responsive yields
Conservation tillage	Residue armour and pore structure preserved	Fuel and labour saving; moisture saving
Agroforestry / grass elements	Hedgerows, bund trees, fodder banks, grass strips	Fodder, fuel and mulch from protective plants

USMAN NOTES www.usmannotes.com

Limitations and Place in the Whole Programme

Agronomic measures have limits that honest teaching must state. They are knowledge- and management-intensive rather than capital-intensive, which means they spread through demonstration and habit change, not through a single construction season, and their full fertility benefits build over several years. On steep slopes and in very erosive climates they cannot by themselves control concentrated runoff, and they presuppose that the farmer controls the land and residue decisions involved — grazing of stubble by other people's

animals, and communal burning customs, can defeat an individual farmer's cover plan unless the community agrees on rules. They are therefore deployed as the first and largest component of an integrated programme, combined with the structural measures of the next question where slope and runoff demand them, and supported by the community organisation of Question 4. Their priority in that programme is justified by simple economics: a rupee spent on cover, rotation and contour working usually prevents more soil loss, over more area, than a rupee spent on structures — and it keeps working, through the farmer's own interest, after the project staff have left.

Conclusion

In conclusion, agronomic measures conserve soil by working through the crop system itself: continuous living or residue cover, contour cultivation, strip cropping, rotations and mixtures that include restorative legumes and leys, cover crops and green manures, fertility management centred on organic matter, conservation tillage, and tree and grass elements woven into the farm. Together they reduce detachment, increase infiltration, trap sediment and rebuild the soil properties on which resistance to erosion depends, while generally raising and stabilising yields. They are the cheapest, widest-reaching and most self-sustaining part of soil conservation, limited mainly on the steepest and wettest land, where they join with structural measures in the integrated treatment this assignment goes on to describe.

Question 3: Structural Measures Used in Erosion Control (20 Marks)

Q. 3 Explain structural measures used in erosion control. (20)

Introduction

Structural measures — also called mechanical or engineering measures — control erosion by physically reshaping the land surface, intercepting and conducting runoff, and stabilising channels and slopes with earth, stone, brushwood or masonry works. They are required where agronomic measures alone cannot cope: on slopes too steep or storms too intense for furrow storage, where runoff is already concentrated into gullies, where field boundaries and roads deliver water onto unprotected land, and where valuable assets such as reservoirs, settlements and valley farmland must be defended. Unlike agronomic measures, they demand surveying and design, significant labour or machinery at construction, land taken out of production for the works themselves, and organised maintenance thereafter — costs that are justified only when the protection they give, and the productive use they make possible, exceed them. This answer explains the principal structural measures in the order a designer meets them on the land: terraces, bunds, diversion and drainage works, waterways and outlets, gully-control structures, and slope and bank stabilisation works, together with the principles of design and maintenance that decide whether structures succeed or fail.

Terraces: Bench, Broad-Base and Graded

Terracing converts a continuous slope into a staircase of shorter, gentler segments, attacking the slope-length and steepness factors directly. The bench terrace, the classical form for steep cultivated hillsides, cuts the slope into level platforms separated by near-vertical risers protected with stone, turf or shrubs; runoff is held on the bench itself, erosion on the treated land is almost eliminated, and intensive crops, orchards and even paddy become possible on

land that was previously destroying itself — at high construction labour, which is economical mainly where land is scarce, valuable and permanently cultivated, as in the hill agriculture of northern Pakistan. Broad-base terraces, used on moderate slopes in mechanised farming, are wide, low ridges and channels that a machine can cross and farm over, dividing the slope without taking strips out of production. Graded terraces are built with a slight longitudinal gradient so that intercepted runoff is conducted slowly to a protected outlet, suiting higher-rainfall areas where ponding the whole storm would overtop the work; level terraces suit drier areas where every millimetre of stored moisture counts. In all types the essentials are the same: correct spacing computed from slope and soil, an adequate channel or bench capacity for the design storm, protected risers and ends, and a safe destination for the water the terrace collects, designed before the terrace lines are marked.

Bunds: Contour, Graded and Field Bunds

Bunding is the most widely used structural measure on cultivated land in South Asia. Contour bunds are small earthen embankments built on the contour across slopes of gentle to moderate gradient, ponding runoff behind them until it infiltrates, conserving both soil and moisture and progressively levelling the inter-bund field as sediment accumulates against the ridge. Graded bunds carry a designed slight slope and discharge surplus water to grassed outlets, suiting heavier rainfall or less permeable soils. Around and between fields, ordinary field bunds — raised, compacted and, where necessary, stone-faced or grass-planted borders — prevent field-to-field flow, hold irrigation and rain where they belong, and mark the stable skeleton on which contour practice is organised; planting bunds with grass, shrubs or trees both strengthens them and yields fodder and fuel from land otherwise unproductive. Bunds fail in predictable ways

that design must pre-empt: built off the contour they drain to a low point and breach there; undersized sections overtop in the design storm; burrowing animals, cart crossings and deliberate cutting by a downhill neighbour open the first gap; and a system of bunds without a planned outlet delivers its collected surplus onto the weakest point of the lowest field. Correct layout on the true contour or grade, adequate cross-section with proper compaction, stone or turf protection of spill points, and agreed maintenance responsibility convert bunding from a frequent disappointment into the workhorse of erosion control.

Diversion Drains, Cutoff Drains and Farm Drainage

Many erosion disasters begin with water arriving from somewhere else — runoff generated on a hillside, a road or a settlement above the field. Diversion drains and cutoff drains intercept that external water before it reaches the land to be protected and convey it, at a non-erosive gradient in a channel of adequate capacity, to a safe outlet such as a grassed waterway, a stable gully or a stream. A cutoff drain above a gully head is often the single most effective gully treatment, because it removes the cause — the plunging inflow — while bed and bank works deal with the accumulated damage. Field and farm drainage works complete the system on flatter land: surface drains that remove excess water without scour, and, in waterlogged and salinising irrigated areas, the drainage whose absence is itself a cause of degradation. Every interception work embodies the governing rule of structural design: never collect water without providing, in advance, a safe place for it to go — a drain that ends on an unprotected slope manufactures a gully more efficiently than nature would have done.

Grassed Waterways, Outlets and Drop Structures

The collected water of a conservation system must descend to the stream somehow, and waterway and outlet structures make that descent safe. Grassed waterways are broad, shallow, parabolic channels in natural drainage lines, kept in dense grass, sized for the peak flow of the contributing area and checked against a permissible velocity for the grass and soil; they receive terrace, bund and drain discharges and deliver them without scour. Where water must fall a significant height — at a terrace outlet, a gully head or a drain exit — energy must be dissipated in a structure rather than in the soil: chute spillways and pipe outlets carry water down a protected path, drop structures step it down in masonry or gabion falls with stilling basins that absorb the plunge, and simple stone or brushwood aprons protect small falls. These outlet works are small in area but decisive in importance, because the failure of a single unprotected outlet can release the concentrated flow of a whole treated slope and unravel the system from its weakest point; designers therefore treat outlets, not the long lengths of bund or terrace, as the critical engineering of the scheme.

Gully Control Structures: Check Dams and Allied Works

Where erosion has concentrated into gullies, structural treatment works in the channel itself. Check dams — barriers built across the gully bed at intervals computed so that the sediment deposited behind one reaches the foot of the next — reduce the effective gradient in steps, trap the moving sediment, raise and stabilise the bed and create the conditions in which vegetation can colonise the banks and floor. According to size, material and intended life they range from brushwood and loose-stone dams for small gullies, through gabion structures of

stone-filled wire baskets that flex and drain, to masonry and concrete dams on large channels; keying into bed and banks, an apron against downstream undercutting, and a spillway section that passes flood flow over the centre of the dam rather than around its ends are the construction details on which survival depends. Supporting works include sloping back and fencing the gully margins, plugging small rills and gully branches with earth and brush, retaining walls where a bank must be held at a road or settlement, and silt-detention and farm ponds that store the intercepted water for use, turning a liability into an irrigation and livestock asset. The structural phase of gully control is deliberately temporary in intent: its purpose is to hold the channel still long enough for the permanent treatment — grass, shrubs and trees — to take over the work.

Slope and Bank Stabilisation Works

The final family of structures protects slopes, banks and disturbed ground directly. Retaining and breast walls of dry stone or masonry hold road cuttings, terrace risers and slide-prone toes where space is too narrow for an earth slope. Gabion spurs and revetments deflect or armour eroding stream banks, and stone pitching protects canal and pond embankments. On landslip areas, surface and subsurface drainage that removes the water lubricating the slide is often more effective than any wall, combined with regrading, benches and deep planting. Contour trenches and staggered contour pits on non-arable slopes — small excavations on the contour that trap runoff and sediment and store moisture for the trees and grass planted with them — bridge the structural and vegetative methods, and contour wattling and fascines of brush laid across small slopes give immediate roughness and trapping while cover establishes. Each such work obeys the general design principles: investigate the cause of instability rather than its

appearance, prefer flexible and drained structures to rigid undrained ones on moving ground, and combine the structure with vegetation from the start.

Table 3: Principal structural measures and their functions

Measure	Function	Typical Setting
Bench / broad-base / graded terraces	Shorten and flatten slope; hold or safely conduct runoff	Steep to moderate cultivated slopes
Contour and graded bunds; field bunds	Pond or conduct runoff; divide slope; retain field water	Gentle to moderate cultivated land, dry to medium rainfall
Diversion / cutoff drains	Intercept external runoff before it reaches protected land	Above fields, gully heads, settlements and roads
Grassed waterways; chutes and drop structures	Convey collected water safely downhill; dissipate energy	Natural drainage lines; all system outlets and falls
Check dams (brush, loose rock, gabion, masonry)	Step down gully gradient; trap sediment; enable revegetation	Rills and gullies of all sizes; ravine reclamation
Retaining walls, spurs, revetments, trenches	Hold or armour slopes and banks; drain unstable ground	Stream banks, road cuttings, landslips, degraded slopes

USMAN NOTES www.usmannotes.com

Design, Construction and Maintenance Principles

Across all these types, three principles decide success. First, design for a stated storm and a stated outlet: structures are sized from the contributing area, slope, soil and a chosen design rainfall, and no work is approved until the destination of its water is fixed and protected. Second, build in the right order and to specification — survey and peg the lines first, construct from the outlet upward and from the ridge downward as the system requires, compact earthworks in layers, key stone work into sound material, and plant the protective vegetation in the same season, not as an afterthought. Third, maintain as a system: annual pre-monsoon inspection, immediate repair of the first small breach — because structures rarely fail whole; they fail at one neglected point and the released water destroys the rest — and a written allocation of responsibility, whether to the landholder, the user group or the agency, agreed before construction begins. Where these principles are observed, structural measures give immediate, measurable protection that agronomy cannot provide alone; where they are ignored, the landscape is littered with breached bunds and undermined dams that discredit conservation itself.

Conclusion

In conclusion, structural measures control erosion by engineering the surface and the drainage: terraces and bunds divide and hold the slope, diversion drains keep foreign water out, grassed waterways and drop structures discharge the collected surplus safely, check dams and allied works arrest and heal gullies, and walls, spurs and trenches stabilise slopes and banks. They are more costly and demanding than agronomic measures and are justified where slope, storm

intensity or existing damage exceeds what cover and cultivation can handle, and they work best — indeed, durably at all — only when built to a design with safe outlets, planted with protective vegetation, and maintained as a permanent system. Structures buy the time and stillness in which biology can recover; the next questions show who must organise that work and how the whole approach is made sustainable.

Question 4: Role of Community Participation in Watershed Management (20 Marks)

Q. 4 Describe the role of community participation in watershed management. (20)

Introduction

A watershed is almost never owned by one person. Its boundary follows the ridge and the stream, crossing the fields of dozens or hundreds of families, common grazing land, public forest, roads and settlements, and the water that one farmer wastes or pollutes on the upper slope becomes the flood or the dry well of another in the valley. Watershed management therefore cannot be delivered to a passive population by an outside agency the way a road or a power line can: nearly every treatment stands on somebody's land, restrains somebody's grazing or requires somebody's maintenance, year after year, long after the project vehicles have left. Community participation — the organised involvement of the people of the watershed in planning, deciding, contributing, implementing, maintaining and monitoring the programme — is consequently not a decorative addition to watershed management but its operating condition. This answer describes the roles participation plays at each stage of the

work, the institutional forms it takes, the benefits it produces, and the conditions and cautions that separate genuine participation from its imitation.

Role in Planning and Decision-Making

Participation begins with planning, and its first role is to supply knowledge no survey team can match and to set priorities the community will actually support. Residents know which spring failed first, where the gully jumped its banks in the great storm, which common land is disputed and which slope the village already protects by custom — the raw material of a realistic watershed plan. Through participatory appraisal methods — village meetings, resource mapping by residents, transect walks across the watershed with farmers and herders, seasonal calendars and problem ranking — the technical team and the community together identify the problems in the order the community feels them, match them to the land's capability, and select treatments that fit local labour, materials and customs. Decisions taken in this way carry a legitimacy that an office-made plan cannot: the choice of which slope to close to grazing, where the check dams go and whose field a waterway must cross has been argued out and accepted in public, with compensation and benefit-sharing settled before the first dispute can harden. Planning participation also corrects the professional's blind spots — a site that is technically ideal for a plantation but is in fact the village's only dry-season grazing, or a drain line that cuts a graveyard — before money is committed to a design that the community would have quietly sabotaged or simply ignored.

Role in Implementation: Labour, Materials and Local Resources

The second role is contribution. Watershed works are labour-hungry — bunds, terraces, trenches, nurseries, planting and check dams over a whole catchment amount to far more work than any project budget can hire — and community labour contribution, whether voluntary, shared by traditional labour-exchange custom, or part-paid under programme rules agreed in advance, is typically what makes treatment of the whole watershed financially possible at all. Communities also contribute local materials — stone, brushwood, grass and tree seed — local transport, and the watchfulness that protects stored material and young works. Contribution has a function beyond cheapness: people maintain what they have built with their own hands and money in a way they never maintain a gift, so the size and willing quality of the local contribution is the best early predictor of whether the works will survive. The terms must, however, be fair and transparent: contributions should be proportionate to benefit and ability, the poorest — who often gain most from fodder, fuel and employment but can least spare unpaid days — should not be driven away by demands designed around the better-off, and accounts of cash and labour should be public, because a single suspicion of misuse can dissolve the cooperation on which everything else depends.

Role in Protection, Maintenance and Rule-Making

The third role is the longest and the most decisive: the daily protection and maintenance of the treated watershed after construction ends. This work is almost entirely a matter of community rules and their enforcement — agreements on closing regenerated slopes to grazing and cutting for an agreed period, rotational grazing arrangements for the pasture, protection of

plantations and waterways from stray animals and fire, regulation of fuelwood cutting, and the routine inspection and repair of bunds, terraces and check dams after each storm season. No department can post a guard on every hillside; only the users themselves, organised in a watershed association, village organisation, water user group or similar body, with rules they have made themselves and sanctions they accept as fair, can supply the continuous low-level enforcement that determines whether cover recovers or is eaten in the first dry season. Experience across the region is remarkably consistent: structurally identical works flourish in villages with functioning user organisations and decay within two or three years in villages where the project treated participation as a formality. Maintenance funds, raised by small regular contributions, fines, nursery income or shares of plantation produce, turn protection from an appeal to goodwill into a small permanent local institution.

Role in Equity, Conflict Resolution and Inclusion

A watershed programme distributes costs and benefits unevenly by nature: upland households accept grazing restrictions and give labour on common slopes, while much of the benefit — steadier water, less flooding, less silt — arrives downhill, and the landless may gain only employment and a share in fodder and fuel. Participation is the mechanism by which these imbalances are recognised and corrected before they destroy the programme: benefit-sharing rules for plantations on common land, priority in employment and in fodder and fuel distribution for those who bore the restrictions, compensation or alternative grazing for households displaced from closed areas, and representation of downstream and upstream hamlets in the same decision body. Participation is also the forum for resolving the resource conflicts — over grazing boundaries, water turns, encroachment and damage by animals — that

degraded watersheds generate constantly, settling them by agreed local process instead of feud or expensive litigation. For participation to perform these roles it must be genuinely inclusive: women, who carry the main burden of fuel, fodder and water and hold much of the practical knowledge of those resources, the landless, tenants and herding groups must have seats and voice, not merely attendance, or the rules produced will protect the interests of the already powerful and fail precisely where enforcement was most needed.

Role in Monitoring, Learning and Continuity

Finally, the community's role extends to watching the results and carrying the work beyond the project. Simple locally kept records — rainfall, well and spring levels, survival of plantations, sediment behind check dams, fodder and fuel yields, repairs done and funds held — allow the village itself to see whether treatment is working, to argue from evidence rather than rumour, and to adapt the rules, for example reopening a protected slope in stages as cover recovers. Local people trained as para-technicians, nursery managers and masons during the programme remain as a permanent skill base in the watershed. When the external agency withdraws, as it always does, it is this combination — an organisation, a fund, a set of rules, local skills and a habit of meeting about the watershed — that constitutes the real product of the project; the bunds and trees are merely its visible first instalment. Participation also disciplines the agency: a community that has contributed and organised will demand the technical support, planting stock and promised funds on time, raising the quality of the professional work itself.

Table 4: Roles of community participation across the life of a watershed programme

Stage	Community Role	What Fails Without It

Planning	Local knowledge, problem ranking, site and treatment decisions, benefit-sharing agreements	Office plans that ignore grazing needs, disputes and customs; works sited wrongly
Implementation	Labour, local materials, transport, supervision of quality	Costs exceed any budget; whole-watershed treatment impossible; no ownership created
Protection and maintenance	Grazing and cutting rules, fire and animal protection, post-storm repair, maintenance fund	Plantations grazed out; first breach unrepaired; works decay in a few seasons
Equity and conflicts	Benefit sharing, compensation, inclusion of women, landless and herders, dispute settlement	Costs fall on upland poor, benefits flow down; resentment destroys cooperation
Monitoring and continuity	Local records, trained local skills, permanent watershed organisation	No learning or adaptation; programme ends with the project

USMAN NOTES www.usmannotes.com

Conditions for Genuine Participation — and Cautions

Participation is not produced by holding meetings. It requires time before construction for trust and organisation to form; honest information about what the programme can and cannot fund; decisions that are actually honoured by the agency when the community chooses within the technical limits; transparency in money and in the selection of office-bearers; and a legal or at least officially recognised standing for the local organisation so that its rules and funds endure. The cautions are the mirror image: participation can be captured by local elites who steer works and wages to themselves; it can degenerate into a ritual of attendance registers used to release funds; women's and herders' nominal presence can conceal their exclusion from the real decisions; and imposed contribution demands can tax the poorest for benefits they receive last. Recognising these risks is not an argument against participation but a specification for doing it properly — the more so because there is no alternative instrument that can protect a watershed at all. Technical quality and participation are complements, not substitutes: the engineer must still design the dam correctly, and the community must still be the body that decides, builds, guards and repairs it.

Conclusion

In conclusion, community participation in watershed management supplies the local knowledge and legitimate decisions that planning requires, the labour and materials that make whole-watershed treatment affordable, the rules, enforcement and maintenance funds on which survival of the works depends, the benefit-sharing and conflict resolution that keep an uneven distribution of costs and benefits fair, and the organisation, skills and records that carry the

work beyond the life of any project. A watershed is a shared resource system, and only the organised community that lives in it can manage it continuously. Programmes that treat people as beneficiaries to be delivered works produce structures that decay; programmes that treat them as the managers, supported by sound technique, produce watersheds that keep improving after the programme closes.

Question 5: Sustainable Land Management Practices (20 Marks)

Q. 5 Explain sustainable land management practices. (20)

Introduction

Sustainable land management (SLM) is the use of land resources — soil, water, vegetation and animals — in ways that meet the changing needs of the present while ensuring the long-term productive potential of those resources and the maintenance of their environmental functions. In other words, it is the positive answer to the land degradation examined in Question 1: where degrading management takes more from the land system than it returns, sustainable management keeps the capital of soil depth, organic matter, cover and water intact while drawing a continuing income of crops, fodder, fuel and other products from it. SLM is not a single technique but an integrated set of practices — agronomic, vegetative, structural and managerial — selected for the capability of each piece of land and combined with the tenure, incentives and community organisation that make them durable. This answer explains the principles of sustainable land management and then describes its main practice groups: soil and fertility management, water management, erosion control and integrated farming systems,

grazing and forest management, and the enabling conditions without which technically sound practices do not persist.

Principles of Sustainable Land Management

Four principles run through all SLM practice. First, use land according to its capability: match the intensity of use to what slope, soil and climate can bear indefinitely, cropping the capable land well, grazing or foresting the fragile land, and protecting the critical land — the matching exercise formalised in land capability classification. Second, keep the soil covered and fed: maintain living or residue cover for as much of the year as possible and return organic matter and nutrients at least equal to removals, because soil organic matter is the working capital from which structure, infiltration and fertility are all drawn. Third, manage water as carefully as soil: capture rain where it falls, store it in the profile and in ponds, dispose of the surplus safely and use every stored unit productively and without generating salinity. Fourth, integrate and diversify: combine crops, livestock, trees and, where relevant, fish and other enterprises so that the wastes of one component feed another, income is spread across seasons and risks, and no single failure can destroy the household. Around these technical principles stand the managerial ones — secure tenure, fair access to common resources, viable economics for each practice in the year it is adopted, not only in the tenth year, and local organisation for the works and rules that exceed one farm — which decide whether practices that are sustainable in the field-trial sense become sustainable in the social sense of actually being continued.

Soil and Fertility Management Practices

The foundation practices work on the soil itself. Organic-matter management — applying farmyard manure and compost, returning crop residues, green manuring, growing legumes in rotation and mixture, and mulching — rebuilds aggregate stability, infiltration, water holding and nutrient supply together, and is the closest thing soil management has to a universal remedy. Balanced and efficient nutrient management replaces what harvests remove, guided where possible by soil testing and by the principle of applying the right source, rate, time and place, combining organic and mineral sources so that fertiliser feeds a biologically active soil rather than substituting for one. Conservation tillage and contour working, described in Question 2, protect this improving soil from physical loss, while biological practices — encouraging earthworms and microbial activity by avoiding unnecessary burning, excessive pesticide use and prolonged bare fallow — maintain the living system that cycles nutrients and builds structure. On problem soils the practices are specific: gypsum, leaching and drainage with salt-tolerant cropping for sodic and saline land, additions of organic matter and careful water management for compacted and crusted surfaces, and agroforestry species chosen for their litter quality where nutrient pumping from depth is needed. The measurable aims are a soil that absorbs the design storm, holds plant-available water through a normal dry spell and yields without a rising input bill — the working definition of a soil being managed sustainably.

Water Management Practices

Sustainable water management on the land begins by harvesting rain in place: contour bunds, terraces, trenches, pits and tied ridges that hold runoff for infiltration; field-level water

harvesting and small farm ponds and tanks that store the managed surplus for livestock and supplementary irrigation at critical crop stages; and recharge structures that direct clean runoff to wells and aquifers. In irrigated systems the corresponding practices are precision land levelling for uniform application, irrigation scheduling that applies water by the crop's need rather than by fixed turns alone, lining or piping of leaky channels where seepage is genuinely lost, alternate and deficit irrigation strategies where research supports them, conjunctive and careful use of groundwater so that pumping does not mine the aquifer or pull up salts, and — indispensably — drainage, surface and subsurface, that removes the excess water and salts which otherwise convert irrigation itself into a degrading force. Watershed-scale water planning links these farm practices: upstream harvesting is sized with regard to downstream users and storages, and riparian buffers and protected catchments guard the quality of the water that all of them share. Done together, these practices raise the productivity of each millimetre of rain and each cubic metre of canal water, which in a water-short country is the largest single production reserve available.

Erosion Control and Integrated Farming Systems

The erosion-control practices of Questions 2 and 3 — cover, contouring, strips and rotations on the agronomic side, bunds, terraces, waterways and check dams on the structural side — form the third group of SLM practices, applied land-unit by land-unit in the ridge-to-valley sequence of a watershed plan. What makes their deployment 'sustainable land management' rather than a list of works is their integration into farming systems that households can profitably operate. Integrated crop-livestock systems recycle manure to fields and residues and fodder to animals, closing the nutrient loop that nutrient mining broke. Agroforestry systems add the tree

component — fodder, fuel, fruit and timber trees on boundaries, bunds, woodlots and contour lines — diversifying income, feeding livestock from protected plots instead of open slopes and building the permanent cover that fragile patches need. Integrated pest and weed management reduces the chemical load on soil and water. Diversification into higher-value, less water-demanding crops where markets allow, and value-adding storage and processing at village level, raise the income earned per unit of soil and water, easing the economic pressure that drives degradation. The test of integration is practical: every enterprise on the farm should either use another's by-product, protect another's resource base, or spread the household's risk — preferably all three.




Figure 2: Illustrative comparison of land outcomes under sustainable land management and under unmanaged, degrading use (index diagram for teaching — not measured data).

Grazing Land, Forest and Common-Land Management

Because a large share of degradation occurs off the cropped field, SLM includes explicit practices for the commons. Rangeland management rests on matching stocking rate to carrying

capacity, rotational and deferred grazing that lets key species seed and recover, protection and reseeding of denuded areas, water-point distribution that spreads grazing pressure, fodder banks and stall-feeding development that carry animals through the dry season without stripping the hills, and community grazing rules of the kind described in Question 4. Sustainable forest and plantation management applies the same logic to trees: harvest not exceeding growth, assisted natural regeneration and enrichment planting, protection from fire and uncontrolled cutting, and a fair local share in forest produce that gives residents an interest in the standing resource. Biodiversity is managed as an asset within all of these — retaining field trees, hedges, wetlands and habitat strips, conserving local crop and livestock varieties, and protecting pollinators and natural enemies — because the ecological services they provide are part of the productive system, not a luxury added to it. Common-land practices fail without the tenure and organisational conditions discussed below, which is why SLM treats commons governance as a practice in its own right rather than a background assumption.

Table 5: Sustainable land management — practice groups and their purposes

Practice Group	Representative Practices	Main Purpose
Soil and fertility	Manure and compost, residues, rotations with legumes, green manuring, balanced nutrients, conservation tillage	Maintain organic matter, structure, infiltration and nutrient capital
Water	In-situ harvesting, ponds and	Capture, store and use water

	recharge, levelling, scheduling, efficient conveyance, drainage	productively without salinity or aquifer mining
Erosion control	Cover, contouring, strips; bunds, terraces, waterways, check dams	Keep soil in place from ridge to valley
Integrated systems	Crop-livestock integration, agroforestry, IPM, diversification	Recycle by-products, spread risk and income, protect fragile patches
Commons management	Stocking control, rotational grazing, regeneration protection, fodder banks	Restore and sustain rangeland, forest and shared resources
Enabling conditions	Secure tenure, incentives, extension, credit, community organisation, monitoring	Make sound practices adoptable, profitable and permanent

USMAN NOTES www.usmannotes.com

Enabling Conditions: Tenure, Incentives, Knowledge and Monitoring

Practices persist only inside enabling conditions, and listing them completes the explanation of SLM. Secure tenure — ownership, long lease or recognised community rights — gives the land user the future from which conservation's delayed benefits are paid, and reform or written

agreements that protect tenants' investments can transform adoption on rented land. Economic incentives align the present with the future: credit suited to multi-year works, fair prices, payments or cost-sharing for the public benefits of conservation such as downstream sediment reduction, and prompt fodder, fuel and income returns from the practices themselves. Knowledge systems — extension, farmer field schools, demonstration on real farms, and the documentation of successful local practice — spread the management skill on which agronomic measures depend. Monitoring closes the loop: simple indicators watched over years — soil organic matter and depth, cover percentage, well levels, sediment behind structures, yield stability — tell managers whether the land is genuinely on a sustainable path or merely busy with projects, and allow correction while correction is still cheap. Policy coherence binds the whole: land, water, forest, livestock and agricultural pricing policies pulling in the same direction, with watershed and land-use planning as the frame in which farm decisions are made.

Conclusion

In conclusion, sustainable land management is the integrated practice of using land within its capability, keeping soil covered and organically fed, capturing and using water without waste or salinity, controlling erosion from ridge to valley, integrating crops, livestock and trees into mutually supporting systems, governing the commons by agreed rules, and sustaining all of it with secure tenure, workable incentives, knowledge and monitoring. Its practices are known, tested and largely affordable; the achievement of the approach is to combine them into systems that a real household, on real land, under a real climate, can continue indefinitely — producing more, more reliably, while handing the soil, water and cover to the next generation

undiminished. That combination of productivity, stability and permanence is the goal toward which every topic in this course — watershed, erosion, conservation measures, structure and participation — has been building.

USMAN NOTES
www.usmannotes.com